

TITLE: Nucleoside syntheses. 19. C-Substitution of nucleosides with the aid of the Eschenmoser sulfide contraction

AUTHOR(S): Vorbruegggen, Helmut; Krolikiewicz, Konrad

CORPORATE SOURCE: Forschungslab., Schering A.-G., Berlin, Fed. Rep. Ger.

SOURCE: Angewandte Chemie (1976), 88(21), 724-5
CODEN: ANCEAD; ISSN: 0044-8249

DOCUMENT TYPE: Journal

LANGUAGE: German

GI For diagram(s), see printed CA Issue.

AB Treatment of thiopurine nucleosides I [$R = SCH_2R_3$ ($R_3 = Bz, Me_3CO_2C, 4-O_2NC_6H_4CH_2$); $R_1 = H, Me_3SiNH$; $R_2 = Ac, Me_3Si$] with strong base and Ph_3P gave C-alkyl nucleosides I ($R = CH_2R_3, R_1 = H, NH_2$) in 72-80% yields. Similarly prepared were II ($X = CH, N; R = CH:C(OH)Ph, R_2 = H$) and III ($R = CH:C(OH)Ph, R_2 = H$) from the corresponding II and III ($R = SCH_2Bz, R_2 = Bz$).

IT 60363-87-3
RL: RCT (Reactant); RACT (Reactant or reagent)
(Eschenmoser sulfide contraction of)

RN 60363-87-3 ZCAPLUS

CN Guanosine, 6-S-(2-oxo-2-phenylethyl)-6-thio-N-(trimethylsilyl)-2',3',5'-tris-O-(trimethylsilyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

